

1 Introduction

Sequences of the form $(\lfloor j\alpha \rfloor)_{j \in \mathbb{N}}$ for $\alpha > 1$, now known as Beatty sequences, were first studied in the context of the famous problem of covering the set of positive integers by disjoint sequences [1]. Further results in the direction of so-called disjoint covering systems are due to [2], [3], [4] and others. Other aspects of Beatty sequences were then studied, such as their generation using graphs [5], their relation to generating functions [6], [7], their substitution invariance [8], [9], etc. A good source of references on Beatty sequences and other related problems can be found in [10], [11].

p03

$$\beta^i = G_i \beta - G_{i-1} \quad \text{and} \quad \beta'^i = G_i \beta' - G_{i-1} \quad (5)$$

The proof we give here is based on the algebraic expression for one-dimensional cut-and-project sets [14]. Let V_1, V_2 be straight lines in \mathbb{R}^2 determined by vectors $(\beta, -1)$ and $(\beta', -1)$, respectively. The projection of the square lattice \mathbb{Z}^2 on the line V_1 along the direction of V_2 is given by

$$(a, b) = (a + b\beta')\bar{x}_1 + (a + b\beta)\bar{x}_2, \text{ for } (a, b) \in \mathbb{Z}^2,$$

where $\bar{x}_1 = \frac{1}{\beta - \beta'}(\beta, -1)$ and $\bar{x}_2 = \frac{1}{\beta' - \beta}(\beta', -1)$. For the de-

scription of the projection of \mathbb{Z}^2 on V_1 it suffices to consider the set

$$\mathbb{Z}[\beta'] := \{a + b\beta' \mid a, b \in \mathbb{Z}\}$$

The integral basis of this free abelian group is $(1, \beta')$, and thus every element x of $\mathbb{Z}[\beta']$ has a unique expression in this base. We will say that a is the rational part of $x = a + b\beta'$ and b is its irrational part. Since β' is a quadratic unit, $\mathbb{Z}[\beta']$ is a ring and, moreover, it satisfies

$$\beta' \mathbb{Z}[\beta'] = \mathbb{Z}[\beta'] \quad (6)$$

A cut-and-project set is the set of projections of points of \mathbb{Z}^2 to V_1 , that are found in a strip of given bounded width, parallel to the straight line V_1 . Formally, for a bounded interval Ω we define

$$\Sigma(\Omega) = \{a + b\beta' \mid a, b \in \mathbb{Z}, a + b\beta \in \Omega\}$$

Note that $a + b\beta'$ corresponds to the projection of the point (a, b) to the straight line V_1 along V_2 , whereas $a + b\beta$ corresponds to the projection of the same lattice point to V_2 along V_1 .

Among the simple properties of the cut-and-project sets that we use here are

$$\Sigma(\Omega - 1) = -1 + \Sigma(\Omega), \quad \beta' \Sigma(\Omega) = \Sigma(\beta\Omega),$$

where the latter is a consequence of (6). If the interval Ω is of unit length, one can derive directly from the definition a simpler expression for $\Sigma(\Omega)$. In particular, we have

$$\Sigma[0, 1) = \{a + b\beta' \mid a + b\beta \in [0, 1)\} = \{b\beta' - \lfloor b\beta \rfloor \mid b \in \mathbb{Z}\}, \quad (7)$$

where we use that the condition $0 \leq a + b\beta < 1$ is satisfied if and only if $a = \lceil -b\beta \rceil = -\lfloor b\beta \rfloor$.

Let us mention that the above properties of one-dimensional cut-and-project sets, and many others, are explained in the review article [14].

3 Self-matching property of the graph $\lfloor j\beta \rfloor$ against j

An important role in the study of the self-matching properties of the graph $\lfloor j\beta \rfloor$ against j is played by the generalized Fibonacci sequence $(G_i)_{i \in \mathbb{N}}$, defined by (2) and (4), respectively. It turns out that shifting the argument j of the function

$\lfloor j\beta \rfloor$ by the integer G_i results in shifting the value by G_{i-1} , with the exception of isolated mismatches with low frequency. The first proposition is an easy consequence of the expressions of β' as an element of the ring $\mathbb{Z}[\beta]$ in the integral basis $1, \beta$, given by (3) and (5).

Theorem 1

Let $\beta \in (0, 1)$ satisfy $\beta^2 + m\beta = 1$ and let $(G_i)_{i=0}^\infty$ be defined by (2). Let $i \in \mathbb{N}$. Then for $j \in \mathbb{Z}$ we have

$$\lfloor \beta(j + G_i) \rfloor = \lfloor j\beta \rfloor + G_{i-1} + \varepsilon_i(j)$$

where $\varepsilon_i(j) \in \{0, (-1)^{i+1}\}$. The frequency of integers j for which the value $\varepsilon_i(j)$ is non-zero is equal to

$$\rho_i := \lim_{n \rightarrow \infty} \frac{\#\{j \in \mathbb{Z} \mid -n \leq j \leq n, \varepsilon_i(j) \neq 0\}}{2n + 1} = \beta^i.$$

Proof. The first statement is trivial. For, we have

$$\begin{aligned} \varepsilon_i(j) &= \lfloor \beta(j + G_i) \rfloor - \lfloor j\beta \rfloor - G_{i-1} = \lfloor j\beta - \lfloor j\beta \rfloor + \beta G_i - G_{i-1} \rfloor \\ &= \lfloor j\beta - \lfloor j\beta \rfloor + (-1)^{i+1} \beta^i \rfloor \in \{0, (-1)^{i+1}\}. \end{aligned} \quad (8)$$

The frequency ρ_i is easily determined in the proof of Theorem 1. \square

In the following theorem we determine the integers j for which $\varepsilon_i(j)$ is non-zero. From this, we easily derive the frequency of such mismatches.

Theorem 2

With the notation of Theorem 1, we have

$$\varepsilon_i(j) = \begin{cases} 0 & \text{if } j \notin U_i, \\ (-1)^{i+1} & \text{otherwise,} \end{cases}$$

where

$$U_i = \{k G_{i+1} + \lfloor k\beta \rfloor G_i \mid k \in \mathbb{Z}, k \neq 0\} \cup \left\{ \frac{(-1)^{i-1}}{2} G_i \right\}.$$

Before starting the proof, let us mention that for i even, the set U_i can be written simply as

$$U_i = \{k G_{i+1} + \lfloor k\beta \rfloor G_i \mid k \in \mathbb{Z}\}.$$

For i odd, the element corresponding to $k = 0$ is equal to $-G_i$ instead of 0. The distinction according to the parity of i is necessary here, since unlike the paper [12], we determine the values of $\varepsilon_i(j)$ for $j \in \mathbb{Z}$, not only for.

Proof. It is convenient to distinguish two cases according to the parity of i .

- First let i be even. It is obvious from (8), that $\varepsilon_i(j) \in \{0, -1\}$ and

$$\varepsilon_i(j) = -1 \quad \text{if and only if} \quad j\beta - \lfloor j\beta \rfloor \in [0, \beta^i). \quad (9)$$

Let us denote by M the set of all such j ,

$$\begin{aligned} M &= \{j \in \mathbb{Z} \mid j\beta - \lfloor j\beta \rfloor \in [0, \beta^i)\} \\ &= \{j \in \mathbb{Z} \mid k + j\beta \in [0, \beta^i), \text{ for some } k \in \mathbb{Z}\} \end{aligned}$$

Therefore M is formed by the irrational parts of the elements of the set

$$\begin{aligned} \{k + j\beta' \mid k + j\beta \in [0, \beta^i)\} &= \Sigma[0, \beta^i) = \beta'^i \Sigma[0, 1) \\ &= (-\beta' G_i + G_{i-1}) \{k\beta' - \lfloor k\beta \rfloor \mid k \in \mathbb{Z}\}, \end{aligned}$$

where the last equality follows from (3) and (7). Separating the irrational part we obtain

$$\begin{aligned} M &= \{k G_i m + k G_{i-1} + \lfloor k\beta \rfloor G_i \mid k \in \mathbb{Z}\} \\ &= \{G_i \lfloor k\beta \rfloor + k G_{i+1} \mid k \in \mathbb{Z}\} = U_i, \end{aligned}$$

where we have used the equations $\beta'^2 + m\beta' = 1$ and $mG_i + G_{i-1} = G_{i+1}$.

- Now let i be odd. Then from (8), $\varepsilon_i(j) \in \{0, -1\}$ and $\varepsilon_i(j) = 1$ if and only if $j\beta - \lfloor j\beta \rfloor \in [1 - \beta^i, 1)$. (10)

Let us denote by M the set of all such j ,

$$\begin{aligned} M &= \{j \in \mathbb{Z} \mid j\beta - \lfloor j\beta \rfloor - 1 \in [-\beta^i, 0)\} \\ &= \{j \in \mathbb{Z} \mid k + j\beta \in [-\beta^i, 0), \text{ for some } k \in \mathbb{Z}\}. \end{aligned}$$

Therefore M is formed by the irrational parts of elements of the set

$$\begin{aligned} \{k + j\beta' \mid k + j\beta \in [-\beta^i, 0)\} &= \Sigma[-\beta^i, 0) = \beta'^i \Sigma[-1, 0) \\ &= \beta'^i (1 - \Sigma[0, 1)) = (\beta' G_i - G_{i-1}) \{k\beta' - \lfloor k\beta \rfloor - 1 \mid k \in \mathbb{Z}\}. \end{aligned}$$

Separating the irrational part we obtain

$$\begin{aligned} M &= \{-k G_i m - k G_{i-1} - \lfloor k\beta \rfloor G_i - G_i \mid k \in \mathbb{Z}\} \\ &= \{-k G_{i+1} - G_i (\lfloor k\beta \rfloor + 1) \mid k \in \mathbb{Z}\} \\ &= \{k G_{i+1} + G_i (\lceil k\beta \rceil - 1) \mid k \in \mathbb{Z}\} = U_i, \end{aligned}$$

where we have used the equation

$$\beta'^2 + m\beta' = 1, \quad mG_i + G_{i-1} = G_{i+1} \quad \text{and} \quad -\lfloor k\beta \rfloor = \lceil k\beta \rceil.$$

Let us recall that the Weyl theorem [15] states that numbers of the form $j\alpha - \lfloor j\alpha \rfloor$, $j \in \mathbb{Z}$, are uniformly distributed in $(0, 1)$ for every irrational α . Therefore the frequency of those $j \in \mathbb{Z}$ that satisfy $j\alpha - \lfloor j\alpha \rfloor \in I \subset (0, 1)$ is equal to the length of the interval I . Therefore one can derive from (9) and (10) that the frequency of mismatches (non-zero values $\varepsilon_i(j)$) is equal to β^i , as stated by Theorem 1. \square

If $\beta \in (0, 1)$ is the quadratic unit satisfying $\beta^2 - m\beta = -1$, then the considerations are even simpler, because expression (5) does not depend on the parity of i . We state the result as the following theorem.

Theorem 3

Let $\beta \in (0, 1)$ satisfy $\beta^2 - m\beta = -1$ and let $(G_i)_{i=0}^\infty$ be defined by (4). For $i \in \mathbb{N}$, put

$$V_i = \{k G_{i+1} - (\lfloor k\beta \rfloor + 1) G_i \mid k \in \mathbb{Z}\}.$$

Then for $j \in \mathbb{Z}$ we have

$$\lfloor \beta(j + G_i) \rfloor = \lfloor j\beta \rfloor + G_{i-1} + \varepsilon_i(j),$$

where

$$\varepsilon_i(j) = \begin{cases} 0 & \text{if } j \notin V_i, \\ 1 & \text{otherwise.} \end{cases}$$

The density of the set U_i of mismatches is equal to β^i .

Proof. The proof follows the same lines as proofs of Theorems 1 and 2. \square

4 Conclusions

One-dimensional cut-and-project sets can be constructed from \mathbb{Z}^2 for every choice of straight lines V_1, V_2 , if they have irrational slopes. However, in our proof of the self-matching properties of the Beatty sequences we strongly use the algebraic ring structure of the set $\mathbb{Z}[\beta']$ and its scaling invariance with the factor β' , namely $\beta' \mathbb{Z}[\beta] = \mathbb{Z}[\beta']$. For this, β' must necessarily be a quadratic unit.

However, it is plausible that, even for other irrationals α , some self-matching property is displayed by the graph $\lfloor j\alpha \rfloor$ against j . To show that, other methods would be necessary.

5 Acknowledgments

The authors acknowledge financial support from the Czech Science Foundation GA R 201/05/0169, and from the grant LC06002 of the Ministry of Education, Youth and Sports of the Czech Republic.

References

- [1] Beatty, S.: *Amer. Math. Monthly*, Vol. 33 (1926), No. 2, p. 103–105.
- [2] Fraenkel, A. S.: The Bracket Function and Complementary Sets of Integers. *Canad. J. Math.*, 21, 1969, 6–27
- [3] Graham, R. L.: Covering the Positive Integers by Disjoint Sets of the Form $\{n\alpha + \beta \mid n = 1, 2, \dots\}$. *J. Combinatorial Theory Ser. A*, Vol. 15 (1973), p. 354–358.
- [4] Tijdeman, R.: Exact Covers of Balanced Sequences and Fraenkel's Conjecture. In *Algebraic Number Theory and Diophantine Analysis* (Graz, 1998), Berlin: de Gruyter 2000, p. 467–483.
- [5] de Bruijn, N. G.: Updown Generation of Beatty Sequences. *Nederl. Akad. Wetensch. Indag. Math.*, Vol. 51 (1989), p. 385–407.
- [6] Komatsu, T.: A Certain Power Series Associated with a Beatty Sequence. *Acta Arith.*, Vol. 76 (1996), p. 109–129.
- [7] O'Bryant, K.: A Generating Function Technique for Beatty Sequences and Other Step Sequences. *J. Number Theory*, Vol. 94 (2002), p. 299–319.
- [8] Komatsu, T.: Substitution Invariant Inhomogeneous Beatty Sequences. *Tokyo Journal Math.*, Vol. 22 (1999), p. 235–243.
- [9] Parvaix, B.: Substitution Invariant Sturmian Bisequences. *Thor. Nombres Bordeaux*, Vol. 11 (1999), p. 201–210.
- [10] Brown, T.: Descriptions of the Characteristic Sequence of an Irrational. *Canad. Math. Bull.*, Vol. 36 (1993), p. 15–21.

- [11] Stolarsky, K.: Beatty Sequences, Continued Fractions, and Certain Shift Operators. *Canad. Math. Bull.*, Vol. **19** (1976), p. 473–482.
- [12] Bunder, M., Tognetti, K.: On the Self Matching Properties of $[j\tau]$. *Discr. Math.*, Vol. **241** (2001), p. 139–151.
- [13] Shechtman, D., Blech, I., Gratias, D., Cahn, J. W.: Metallic Phase with Long-Range Orientational Order and no Translation Symmetry. *Phys. Rev. Lett.*, Vol. **53** (1984), p. 1951–1953.
- [14] Gazeau, J. P., Masáková, Z., Pelantová, E.: Nested Quasicrystalline Discretization of the Line. In: *Physics and Number Theory* (Editor: L. Nyssen), Vol. 10 of *IRMA Lectures in Mathematics and Theoretical Physics*, Zürich, EMS 2006, p. 79–132.
- [15] Weyl, H.: Über die Gleichverteilung von Zahlen mod. Eins. *Math. Ann.*, Vol. **77** (1916), p. 313–352.

Doc. Ing. Zuzana Masáková, Ph.D.
phone: + 420 224 358 544
e-mail: masakova@km1.fjfi.cvut.cz,

Prof. Ing. Edita Pelantová, CSc.
phone: + 420 224 358 544
e-mail: pelantova@km1.fjfi.cvut.cz

Doppler Institute for Mathematical Physics and Applied Mathematics

Czech Technical University in Prague
Faculty of Nuclear Sciences and Physical Engineering
Trojanova 13
120 00 Praha 2, Czech Republic